



DynaMed[®]

Nuevas funcionalidades:
Drug Interaction checker y Drugs A-Z

Opciones de consulta de fármacos

DynaMed

EP SALES
SIGN IN

Learn more about CME

Specialties ▾

Drugs A-Z

Drug Interactions

Differential Diagnosis

Calculators

About

Mobile

CONFIDENCE IN PRACTICE


🌐

Español

▾

🔍

Buscar



Cardiology

Acute Coronary Syndromes,
Stable Coronary Artery Disease,
Heart Failure

View Specialty



Oncology

Breast Cancers, Lung Cancers,
Gastrointestinal Cancers,
Lymphomas

View Specialty



Dermatology

Eczematous Dermatoses,
Malignant and Pre-malignant
Lesions, Papulosquamous...

View Specialty

Feedback

Opciones de consulta: Drugs A-Z

DynaMed



EP SALES
SIGN IN

Learn more about CME



Buscar

Specialties ▾ Drugs A-Z Drug Interactions Differential Diagnosis Calculators About Mobile

Drugs A-Z

Powered by *IBM Micromedex*



A

B

C

D

E

F

G

H

I

J

K

L

M

N

O

P

Q

R

S

T

U

V

>

Fex

F

Fexofenadine



Fexofenadine/Pseudoephedrine

Solo hay que pinchar sobre la primera letra y luego utilizar la caja de búsqueda o el listado para localizar el fármaco que queremos buscar



Acceda rápidamente a
cada una de las
secciones dentro del tema

Fexofenadine

IBM Micromedex®

Class

Class

- > Dosing & Indications
- > Contraindications/Warnings
- Pregnancy & Lactation
- Drug Interactions
- > Adverse Effects
- Monitoring
- > Toxicology
- Clinical Teaching
- Administration
- Mechanism of Action
- Pharmacokinetics
- US Availability
- > How Supplied
- References

Class

- Antihistamine, Less-Sedating
- Butyrophenone
- Respiratory Agent

Dosing & Indications

Adult Dosing

- Idiopathic urticaria, chronic
 - 60 mg ORALLY twice daily OR 180 mg ORALLY once daily with water (manufacturer dosing) ; if symptoms persist after 2 weeks, may increase up to 4 times the standard dose (guideline dosing)
- Seasonal allergic rhinitis
 - 60 mg orally twice daily or 180 mg once daily with water
- [IBM Micromedex® DRUGDEX® Subscribers: FEXOFENADINE Details](#)

Pediatric Dosing

- General Dosage Information
 - safety and efficacy not established in children less than 6 months of age
 - orally disintegrating tablets (ODT) not approved for children under 6 years of age

Opciones de consulta: Drug interaction checker

DynaMed



EP SALES
SIGN IN

[Learn more about CME](#)



Buscar

[Specialties](#) ▾

[Drugs A-Z](#)

[Drug Interactions](#)

[Differential Diagnosis](#)

[Calculators](#)

[About](#)

[Mobile](#)

Drug Interactions

Use the search fields to look up two or more drugs for potential interactions.

SIMVASTATIN



FEXOFENADINE



ZOLPIDEM



CLOPIDOGREL



ASPIRIN



[+](#) Add Drug

Check for Interactions

[Start Over](#)

Elija los fármacos para, de forma rápida, chequear posibilidades de interacciones entre ellos

Interactions for SIMVASTATIN, FEXOFENADINE, ZOLPIDEM, CLOPIDOGREL, ASPIRIN

Display

Drug/Drug interactions

Severity Index

Drug/Drug Interactions (2):

CLOPIDOGREL HYDROGEN SULFATE

MAJOR

Documentation: Fair

Concurrent use of **ASPIRIN** and **CLOPIDOGREL** may result in an increased risk of bleeding. [See details >](#)


SIMVASTATIN

MODERATE

Documentation: Excellent

Concurrent use of **CLOPIDOGREL** and **CYP3A4 METABOLIZED STATINS** may result in decreased formation of clopidogrel active metabolite resulting in high on-treatment platelet reactivity. [See details >](#)

Acceso disponible 24h/día en cualquier parte del mundo



DynaMed®

Benedita Menezes

bmenezes@ebSCO.com

+34 638 931 755